

Fibrous dysplasia of the spine with sarcomatous transformation: a case report and review of the literature

E. Yalniz, T. Er, and F. Özyilmaz

Department of Orthopaedic Surgery, Faculty of Medicine, University of Trakya, Edirne, Turkey

Summary. A fibrosarcoma is reported in the spine of a 53-year-old man with polyostotic fibrous dysplasia. There was no history of endocrine disturbances and no previous irradiation. Malignant transformation in fibrous dysplasia is rare. A review of the literature reveals 101 cases of malignant degeneration occurring in fibrous dysplasia. We believe that this is the first report of sarcomatous change arising in an area of fibrous dysplasia in the spine.

Key words: Fibrous dysplasia – Fibrosarcoma – Malignant transformation

Fibrous dysplasia is a common benign pathological condition characterized by fibroosseous metaplasia. There are monostotic and polyostotic forms. Polyostotic fibrous dysplasia may be accompanied by skin pigmentation, endocrine disorders, and precocious puberty, an entity known as Albright's syndrome [1].

Although any bone can be affected, vertebral column involvement is uncommon, especially in the monostotic type [7, 16, 18, 20]. Thoracic location is even more unusual. Wright and Stoker [20] reported nine cases of vertebral involvement in polyostotic fibrous dysplasia; in only one of these cases was there a lesion in the thoracic region.

Malignant degeneration of fibrous dysplasia is a rare phenomenon. The most common malignant tumor is osteosarcoma (53.4%) followed by fibrosarcoma (17.8%) and chondrosarcoma (8.9%).

Case report

A 53-year-old man complained of back pain and weakness. His blood count, ESR, and alkaline phosphatase level were normal. No previous radiotherapy had been administered. Plain roentgenograms showed expanding lytic lesions involving the thoracolumbar spine

and multiple ribs. Computerized axial tomography (CT) and magnetic resonance imaging (MRI) revealed a compressed spinal cord between T7 and T10 (Fig. 1). Biopsy of the left tenth rib revealed typical histopathological changes in fibrous dysplasia. The patient underwent an exploration of his thoracic and lumbar spine through a posterior approach. A decompressive laminectomy between T7 and T11 was done with transpedicular vertebral biopsy, and posterior spinal instrumentation was applied. Histological examination of the specimen from the vertebral body and posterior elements revealed fibrous dysplasia with features of fibrosarcoma (Fig. 2). The patient's symptoms were relieved and his motor function improved. He then received six courses of chemotherapy (Adriamycin). Twenty-eight months after the operation, he was still alive (Fig. 3).

Discussion

Fibrous dysplasia, in either monostotic or polyostotic form, rarely involves the vertebral column. Campanacci [3] reviewed 166 patients and found 2 with monostotic and 2 with polyostotic fibrous dysplasia of the spine. Dahlin and Unni [5] studied 471 patients who had fibrous dysplasia; only 6 patients had vertebral involvement. Schajowicz [14] reported 222 monostotic cases and 36 polyostotic cases, including a sacral lesion in the monostotic group. There is no predilection for any particular portion of the spine. Both the posterior elements and vertebral bodies are affected. When a lesion of the vertebral body becomes extensive, vertebral collapse may occur. Cord compression or nerve encroachment may be caused by expansion of either the vertebral body itself or of posterior elements.

Malignant degeneration of fibrous dysplasia is a rare but well-recognized complication of which 101 cases have been reported [2, 4, 6, 8–13, 15, 17, 19, 21]. The incidence is estimated at 0.4% (6/1517) for fibrous dysplasia and 4% for Albright's syndrome [15]. The common sites were craniofacial bones (35.6%), femur (24.7%), and tibia (12.8%). There has been no other case report regarding sarcomatous transformation in fibrous dysplasia of the spine.

Of the reported cases, 44 patients had monostotic fibrous dysplasia, 46 the polyostotic form, and 11 had Albright's syndrome. There was no predilection for either sex. Osteosarcoma was the most common type of sarcoma that developed in fibrous dysplasia (54 cases). The next most common tumors were fibrosarcoma (18 cases) and

Received: 13 May 1994

Revised: 12 September 1994

Accepted: 28 September 1994

Correspondence to: E. Yalniz, Department of Orthopaedic Surgery, Faculty of Medicine, University of Trakya, 22030 Edirne, Turkey





Fig. 1. **A, B** Anteroposterior and lateral roentgenograms of the thoracolumbar spine show expanding lytic irregular lesions, collapse of the vertebral bodies, and kyphosis. **C** Computerized axial tomogram of the seventh thoracic vertebra. Note the involvement of the body, pedicle, lamina, and the right rib. **D** Magnetic resonance imaging demonstrates compressed spinal cord between T7 and T10

Fig. 2. **A** Typical histological appearance of fibrous dysplasia. Irregular claw-shaped trabeculae of bone are separated by fibrous connective tissue. Hematoxylin-eosin stain, $\times 80$. **B** Histological appearance of fibrosarcoma. Hematoxylin-eosin stain, $\times 150$

Fig. 3A, B. Anteroposterior and lateral roentgenograms taken 28 months after surgery

chondrosarcoma (9 cases). Ebata et al. [6] reported a case of polyostotic fibrous dysplasia in which two types of malignant tumor arose: a chondrosarcoma and an osteosarcoma. Malignant degeneration usually develops in the third or fourth decade of life. The average age at onset is 32 years, and the average lag between the development of fibrous dysplasia and sarcoma is 13.5 years [15].

The role of irradiation in malignant degeneration of fibrous dysplasia has been discussed. In the review by Yabut et al. [21], 23 of 83 patients had undergone prior radiation therapy. Chetty et al. [4] reported three cases of malignant neoplasms occurring in fibrous dysplasia in-

volving facial bones. Two of the tumors had been irradiated. The mean interval between radiation therapy and the onset of the sarcoma was 11 years [15]. In this study, 28 patients (27.7%) had a history of irradiation as treatment for fibrous dysplasia. In 20 of them (71.4%), the tumor was an osteosarcoma. It seems that irradiation provokes the fibrous dysplasia to undergo sarcomatous change. Therefore, radiotherapy should not be used for the treatment of fibrous dysplasia.

The case described here is of fibrosarcoma arising in fibrous dysplasia of the vertebral column without prior irradiation. The location of the disease is very exceptional.

As the patient was considered unfit for radical resection of the tumor, he was treated only with posterior decompression and chemotherapy, with good palliative results.

References

1. Albright F, Butter AM, Smith P (1937) Syndrome characterized by osteitis fibrosa disseminata, areas of pigmentation and endocrine dysfunction, with precocious puberty in females: report of five cases. *N Engl J Med* 216:727-746
2. Blackwell JB (1993) Mesenchymal chondrosarcoma arising in fibrous dysplasia of the femur. *J Clin Pathol* 46:961-962
3. Campanacci M (1990) Bone and soft tissue tumors. Springer, Vienna New York, pp 391-417
4. Chetty R, Kalan MR, Kranold DH (1990) Malignant transformation in fibrous dysplasia. A report of 3 cases. *S Afr J Surg* 28:80-82
5. Dahlin DC, Unni KK (1986) Conditions that commonly simulate primary neoplasms of bone. In: Dahlin DC, Unni KK (eds) Bone tumors. General aspects and data on 8542 cases, 4th edn. Thomas, Springfield, Ill, pp 413-420
6. Ebata K, Usami T, Tohrai I, Kaneda T (1992) Chondrosarcoma and osteosarcoma arising in polyostotic fibrous dysplasia. *J Oral Maxillofac Surg* 50:761-764
7. Ehara S, Kattapuram SV, Rosenberg AE (1992) Fibrous dysplasia of the spine. *Spine* 17:977-979
8. Ishida T, Machinami R, Kojima T, Kikuchi F (1992) Malignant fibrous histiocytoma and osteosarcoma in association with fibrous dysplasia of bone. Report of three cases. *Pathol Res Pract* 188:757-763
9. Jose CC, Benjamin CS (1986) Osteogenic sarcoma arising in polyostotic fibrous dysplasia. A case report. *Australas Radiol* 30:134-136
10. Mock D, Rosen IB (1986) Osteosarcoma in irradiated fibrous dysplasia. *J Oral Pathol* 15:1-4
11. Mortensen A, Bojsen-Moller M, Rasmussen P (1989) Fibrous dysplasia of the skull with acromegaly and sarcomatous transformation. Two cases with a review of the literature. *J Neurooncol* 7:25-29
12. Present D, Bertoni F, Enneking WF (1986) Osteosarcoma of the mandible arising in fibrous dysplasia. A case report. *Clin Orthop* 204:238-244
13. Roth A, Touret P, Rigault P (1985) Association of Ewing's sarcoma with fibrous dysplasia of the tibia. Report of one case. *Rev Chir Orthop Reparatrice Appar Mot* 71:133-137
14. Schajowicz F (1981) Tumor and Tumorlike Lesions of Bone Joints. Springer, New York Berlin Heidelberg, pp 478-490
15. Schwartz DT, Alpert M (1964) The malignant transformation of fibrous dysplasia. *Am J Med Sci* 247:1-20
16. Serena S, Healey JH, Huvos AG (1990) Fibrous dysplasia of the second cervical vertebra. *J Bone Joint Surg [Am]* 72:781-793
17. Taconis WK (1988) Osteosarcoma in fibrous dysplasia. *Skeletal Radiol* 17:163-170
18. Troop JK, Herring JA (1988) Monostotic fibrous dysplasia of the lumbar spine. Case report and review of the literature. *J Pediatr Orthop* 8:599-601
19. Witkin GB, Guilford WB, Siegal GP (1986) Osteogenic sarcoma and soft tissue myxoma in a patient with fibrous dysplasia and hemoglobins. *Clin Orthop* 204:245-252
20. Wright JFS, Stoker DJ (1988) Fibrous dysplasia of the spine. *Clin Radiol* 39:523-527
21. Yabut SM, Kenan S, Sissons HA, Lewis MM (1988) Malignant transformation of fibrous dysplasia. A case report and review of the literature. *Clin Orthop* 228:281-289